


## New Onset Adult Microscopic Hematuria Guideline

### Guideline History

Date Approved	03/00, 02/04, 01/06
Date Revised	01/02, 01/04, 01/06
Date Reviewed	10/05
Next Review Date	01/08

These Guidelines are promulgated by Sentara Healthcare (SHC) as recommendations for the clinical management of specific conditions. Clinical data in a particular case may necessitate or permit deviation from these Guidelines. The SHC Guidelines are institutionally endorsed recommendations and are not intended as a substitute for clinical judgment.

## *New Onset Adult Microscopic Hematuria*



**NOTE:** Asymptomatic gross hematuria needs automatic referral to Urology, Nephrology or Radiology for consult.

**Recommendation:** If patients have other risk factors, i.e. Hx of heavy smoking, urothelial malignancies in the families or others they may need to be screened with upper tract studies and cystourethroscopic evaluations on a first visit rather than waiting 6 weeks to 3 months for next visit.

These guidelines are promulgated by Sentara Healthcare (SHC) as recommendations for the clinical management of specific conditions. Clinical data in a particular case may necessitate or permit deviation from these guidelines. The SHC guidelines are institutionally endorsed recommendations and are not intended as a substitute for clinical judgment.

Approved: 03/00, 02/04, 01/06  
Revised: 01/02, 01/04, 01/06  
Reviewed: 10/05

## **REFERENCES**

### *Urology*

#### **Microscopic Hematuria**

Ostroff, Edward. Personal Communication. October 2005.