



Chronic Low Back Pain Guideline

For complete guideline, please go to www.aafp.org

or

<http://www.aafp.org/patient-care/clinical-recommendations/all/back-pain.html>

Guideline History

Date Approved	05/03
Review/Revised	10/03, 10/05, 10/07, 11/09, 3/10, 10/12, 9/13, 9/15, 9/17
Next Review Date	9/19

Chronic Low Back Pain: Evaluation and Management

ALLEN R. LAST, MD, MPH, and KAREN HULBERT, MD, *Racine Family Medicine Residency Program, Medical College of Wisconsin, Racine, Wisconsin*

Chronic low back pain is a common problem in primary care. A history and physical examination should place patients into one of several categories: (1) nonspecific low back pain; (2) back pain associated with radiculopathy or spinal stenosis; (3) back pain referred from a nonspinal source; or (4) back pain associated with another specific spinal cause. For patients who have back pain associated with radiculopathy, spinal stenosis, or another specific spinal cause, magnetic resonance imaging or computed tomography may establish the diagnosis and guide management. Because evidence of improved outcomes is lacking, lumbar spine radiography should be delayed for at least one to two months in patients with nonspecific pain. Acetaminophen and nonsteroidal anti-inflammatory drugs are first-line medications for chronic low back pain. Tramadol, opioids, and other adjunctive medications may benefit some patients who do not respond to nonsteroidal anti-inflammatory drugs. Acupuncture, exercise therapy, multidisciplinary rehabilitation programs, massage, behavior therapy, and spinal manipulation are effective in certain clinical situations. Patients with radicular symptoms may benefit from epidural steroid injections, but studies have produced mixed results. Most patients with chronic low back pain will not benefit from surgery. A surgical evaluation may be considered for select patients with functional disabilities or refractory pain despite multiple nonsurgical treatments. (*Am Fam Physician*. 2009;79(12):1067-1074. Copyright © 2009 American Academy of Family Physicians.)

► **Patient information:** A handout on coping with chronic low back pain, written by the authors of this article, is available at <http://www.aafp.org/afp/20090615/1067-s1.html>.



This clinical content conforms to AAFP criteria for evidence-based continuing medical education (EB CME).

Most primary care physicians can expect to see at least one patient with low back pain per week. Acute episodes of back pain are usually self-limited. Patients with persistent or fluctuating pain that lasts longer than three months are defined as having chronic low back pain. Patients with chronic low back pain are more likely to see a family physician (65.0 percent) for their pain compared with orthopedists (55.9 percent), physical therapists (50.5 percent), and chiropractors (46.7 percent).¹ The economic impact of chronic low back pain stems from prolonged loss of function, resulting in loss of work productivity, treatment costs, and disability payments. Estimates of these costs range from \$12.2 to \$90.6 billion annually.¹

Evaluation

The initial evaluation, including a history and physical examination, of patients with chronic low back pain should attempt to place patients

into one of the following categories: (1) nonspecific low back pain; (2) back pain associated with radiculopathy or spinal stenosis; (3) back pain referred from a nonspinal source; or (4) back pain associated with another specific spinal cause² (*Table 1*³). For patients who have back pain associated with radiculopathy, spinal stenosis, or another specific spinal cause, magnetic resonance imaging (MRI) or computed tomography (CT) may establish the diagnosis and guide management.

The medical history should include questions about osteoporosis, osteoarthritis, and cancer, and a review of any prior imaging studies. Review of systems should focus on unexplained fevers, weight loss, morning stiffness, gynecologic symptoms, and urinary and gastrointestinal problems.

The physical examination should include the straight leg raise and a focused neuromuscular examination. A positive straight leg raise test (pain with the leg fully extended at the knee and flexed at the hip between

Table 1. Differential Diagnosis of Chronic Low Back Pain

Nonspecific or idiopathic (70 percent)	Referred pain (2 percent)	Nonmechanical (1 percent)
Lumbar sprain or strain	Aortic aneurysm	Neoplasia
Mechanical (27 percent)	Diseases of the pelvic organs	Multiple myeloma
Degenerative processes of disks and facets	Prostatitis	Metastatic carcinoma
Herniated disk	Endometriosis	Lymphoma and leukemia
Osteoporotic fracture*	Chronic pelvic inflammatory disease	Spinal cord tumors
Spinal stenosis	Gastrointestinal disease	Retroperitoneal tumors
Traumatic fracture*	Pancreatitis	Primary vertebral tumors
Congenital disease	Cholecystitis	Inflammatory arthritis, often associated with human leukocyte antigen-B27
Severe kyphosis	Penetrating ulcer	Ankylosing spondylitis
Severe scoliosis	Renal disease	Psoriatic spondylitis
Transitional vertebrae	Nephrolithiasis	Reiter syndrome
Spondylosis	Pyelonephritis*	Inflammatory bowel disease
Internal disk disruption or discogenic pain	Perinephric abscess*	Infection*
Presumed instability		Osteomyelitis
		Septic diskitis
		Paraspinal abscess
		Epidural abscess
		Shingles
		Scheuermann disease (osteocondrosis)
		Paget disease of bone

*—Indicates conditions more likely to present as acute low back pain.

Adapted with permission from Deyo RA, Weinstein JN. Low back pain. *N Engl J Med*. 2001;344(5):365.

rapidly progressive disease (Table 2^{5,6}) or radicular symptoms that do not spontaneously resolve after six weeks. Because evidence of improved outcomes is lacking, imaging, such as lumbar spine radiography, should be delayed at least one to two months in patients with nonspecific pain without red flags for serious disease.⁶

Psychosocial issues play an important role in guiding the treatment of patients with chronic low back pain. One study found that patients with chronic low back pain who have a reduced sense of life control, disturbed mood, negative self-efficacy, high anxiety levels, and mental health disorders, and who engage in catastrophizing tend to not respond well to treatments such as epidural steroid injections.⁸ “Yellow flags” are psychosocial risk factors for long-term disability⁹ (Table 3⁹⁻¹¹). Evaluation of psychosocial problems and “yellow flags” are useful in identifying patients with a poor prognosis.^{8,9}

Management

GENERAL PRINCIPLES

The goals of treating chronic low back pain often change over time, shifting from the

initial intent to cure to improving pain and function. Patients often have unrealistic expectations of complete pain relief and full return to their previous level of activity. There is often a large gap between a patient’s desired amount of pain reduction and the minimum percentage of improvement that would make a treatment worthwhile.¹² Documenting goals and expectations and revisiting them on follow-up visits may be helpful.

Patients should receive information about effective self-care options and should be advised to remain active (because muscles that do not move can eventually become hypersensitive to pain).¹³ Assessing the response to therapy should focus on improvements in pain, mood, and function.

Treatment should begin with maximal recommended doses of nonsteroidal anti-inflammatory drugs (NSAIDs) and acetaminophen, followed by adjunctive medications. Nonpharmacologic therapies are effective in certain clinical situations and can be added to the treatment program at any time. For those with severe functional disabilities, radicular symptoms, or refractory pain, referral for epidural steroid injection or surgical evaluation may be reasonable (Figure 1²).

30 and 70 degrees) can suggest lumbar disk herniation, with ipsilateral pain being more sensitive (i.e., better at ruling out disk herniation if negative) and contralateral pain being more specific (i.e., better at ruling in herniation if positive).⁴ Testing deep tendon reflexes, strength, and sensation can help identify which nerve roots are involved.

Laboratory assessment, including erythrocyte sedimentation rate, complete blood count, and C-reactive protein level, should be considered when red flags indicating the possibility of a serious underlying condition are present (Table 2^{5,6}). Urinalysis may be useful when urinary tract infections are suspected, and alkaline phosphatase and calcium levels can help identify conditions, such as Paget disease of bone, that affect bone metabolism; however, these tests are not needed in all patients with chronic low back pain.

Imaging has limited utility because most patients with chronic low back pain have nonspecific findings on imaging studies,⁷ and asymptomatic patients often have abnormal findings.⁶ Initial imaging with MRI, which is the preferred study, or CT is only recommended for patients with red flags for serious or

Table 2. Red Flags Indicating Serious Causes of Chronic Low Back Pain and Evaluation Strategies

Finding	Diagnosis of concern				Evaluation strategy		
	Cauda equina syndrome	Fracture	Cancer	Infection	CBC/ESR/CRP level	Plain radiography	MRI
Age older than 50 years		X	X		1*	1	2
Fever; chills; recent urinary tract or skin infection; penetrating wound near spine				X	1	1	1
Significant trauma		X				1	2
Unrelenting night pain or pain at rest			X	X	1*	1	2
Progressive motor or sensory deficit	X		X				1E
Saddle anesthesia; bilateral sciatica or leg weakness; difficulty urinating; fecal incontinence	X						1E
Unexplained weight loss			X		1*	1	2
History of cancer or strong suspicion for current cancer			X		1*	1	2
History of osteoporosis		X				1	2
Immunosuppression				X	1	1	2
Chronic oral steroid use		X		X	1	1	2
Intravenous drug use		X		X	1	1	2
Substance abuse		X		X	1	1	2
Failure to improve after six weeks of conservative therapy			X	X	1*	1	2 (or unnecessary)

NOTE: Red flags indicate the possibility of a serious underlying condition.

1 = first-line evaluation in most situations; 2 = follow-up evaluation; CBC = complete blood count; CRP = C-reactive protein; E = emergent evaluation required; ESR = erythrocyte sedimentation rate; MRI = magnetic resonance imaging.

*—Prostate-specific antigen testing may be indicated in men in whom cancer is suspected.

Adapted from Kinkade S. Evaluation and treatment of acute low back pain. *Am Fam Physician.* 2007;75(8):1184, with additional information from reference 6.

Table 3. Psychosocial “Yellow Flags” Predicting Long-Term Disability in Patients with Chronic Low Back Pain**Affect**

Anxiety; depression; feeling of uselessness; irritability

Behavior

Adverse coping strategies; impaired sleep because of pain; passive attitude about treatment; withdrawal from activities

Beliefs

Thinks “the worst” or that pain is harmful or uncontrollable, or that it needs to be eliminated (before returning to activities or work)

Social

History of sexual abuse, physical abuse, or substance abuse; lack of support; older age; overprotective family

Work

Expectation that pain will increase with work and activity; pending litigation; problems with worker’s compensation or claims; poor job satisfaction; unsupportive work environment

Information from references 9 through 11.

PHARMACOLOGIC TREATMENT OPTIONS

Acetaminophen is first-line therapy because of its high safety profile. NSAIDs provide similar analgesia, but have significant gastrointestinal and renovascular adverse effects.^{2,14} There are several classes of NSAIDs, and if one class fails, medications from other classes can be tried before abandoning them altogether (*Table 4*). Tramadol (Ultram), opioids, and other adjunctive medications may benefit some patients who do not respond to NSAIDs.

Tramadol is an analgesic that has weak opioid and serotonin-norepinephrine reuptake inhibitor (SNRI) activity. Studies demonstrate short-term improvements in pain and function, but long-term data are lacking.^{15,16} Because of its opioid activity, tramadol generally should not be used in patients recovering from narcotic

Treatment of Chronic Low Back Pain




Figure 1. Treatment algorithm for patients with chronic low back pain.

Adapted with permission from Chou R, Qaseem A, Snow V, et al., for the Clinical Efficacy Assessment Subcommittee of the American College of Physicians, American Pain Society Low Back Pain Guidelines Panel. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society [published correction appears in *Ann Intern Med.* 2008;148(3):247-248]. *Ann Intern Med.* 2007;147(7):482.

addiction. Adverse effects include drowsiness, constipation, and nausea.

All muscle relaxants provide similar short-term improvements in pain and function, but there is no evidence to support their long-term use for chronic low back pain. Sedation is a common adverse effect, and chronic use of benzodiazepines and carisoprodol (Soma) carries the risk of dependency.¹⁷

A 2006 Cochrane review¹⁸ found that some herbal medications appear effective in short-term randomized trials, but lack long-term safety data. *Harpagophytum procumbens* (devil's claw) in a dosage of 50 mg daily, *Salix alba* (white willow bark, a source of salicylic acid) in a dosage of 240 mg daily, and *Cap-sicum frutescens* (cayenne) plaster applied topically every day appear to be better than placebo at reducing chronic low back pain. Limited studies have shown that devil's claw and white willow bark appear to be as effective as NSAIDs.¹⁸

Short-acting (immediate-release) and long-acting (sustained-release) opioid analgesics are sometimes used for chronic low back pain. There have been few high-quality trials to assess the effectiveness and potential risks of these medications in chronic low back pain.¹⁹

Taking opioids can lead to the development of tolerance, hyperalgesia (enhanced pain response to noxious stimuli), and allodynia (enhanced pain response to innocuous stimuli).²⁰ The combination of tolerance and hyperalgesia can decrease opioid effectiveness over time. One of the challenges of treating chronic low back pain is differentiating among tolerance, opioid-induced hyperalgesia, and disease progression. Hyperalgesia involves increasing pain despite increasing opioid treatment, pain that becomes more diffuse and beyond the distribution of the preexisting pain, and an apparent change in pain threshold or tolerability.²⁰ In this situation, the dosage of opioids should be decreased, or patients should be weaned off the medication altogether.

Selective serotonin reuptake inhibitors, SNRIs, and antiepileptic medications have not been shown to help patients with chronic low back pain. Tricyclic antidepressants, however, provide some benefit and can be a

Table 4. Classes of Nonsteroidal Anti-inflammatory Drugs for Chronic Low Back Pain

Class	Generic (brand)	Standard dosage	Maximal dosage (mg per day)	Approximate monthly cost*
Salicylic acids	Aspirin	325 to 650 mg every four hours	4,000	\$3 for 325-mg dose
	Diflunisal (Dolobid)	500 mg two times daily	1,500	\$77 (generic) and \$73 (brand)
	Salsalate	1,500 mg two times daily	3,000	\$27 to \$40
	Choline magnesium trisalicylate	1,500 mg two times daily	3,000	\$44 to \$54
Acetic acids	Diclofenac potassium (Cataflam)	50 mg three times daily	200	\$140 to \$173 (generic) and \$327 (brand)
	Diclofenac sodium, delayed release (Voltaren)	50 mg two or three times daily	200	\$85 to \$98 (generic) and \$192 (brand) for 50 mg two times daily
	Etodolac	200 to 400 mg two or three times daily	1,200	\$77 to \$90 for 200 mg two times daily
	Indomethacin (Indocin)	25 to 50 mg three times daily	200	\$5 to \$30 (generic) and \$80 (brand) for 25-mg dose
	Indomethacin, extended release (Indocin SR)	25 to 50 mg one or two times daily	150	\$60 (generic) and \$84 (brand) for 25 mg once daily
	Sulindac (Clinoril)	200 mg two times daily	400	\$72 to \$80 (generic) and \$86 (brand)
	Tolmetin	200 to 600 mg three times daily	1,800	\$67 for 200-mg dose
Oxicam	Meloxicam (Mobic)	7.5 to 15 mg once daily	15	\$95 to \$108 (generic) and \$117 (brand) for 7.5-mg dose
	Piroxicam (Feldene)	20 mg once daily	20	\$79 to \$104 (generic) and \$133 (brand)
Propionic acids	Ibuprofen	600 mg four times daily or 800 mg three times daily	2,400	\$30 to \$35 (generic) and \$48 for 600-mg dose
	Ketoprofen	50 to 100 mg three times daily	300	\$60 to \$204 for 50-mg dose
	Naproxen (Naprosyn)	250 to 500 mg two times daily	1,500	\$42 to \$72 (generic) and \$70 (brand) for 250-mg dose
	Naproxen sodium (Anaprox)	275 to 550 mg two times daily	1,650	\$50 to \$53 (generic) and \$63 (brand) for 275-mg dose
	Oxaprozin (Daypro)	1,200 mg once daily	1,800	\$108 to \$164 (generic) and \$157 (brand)
Anthranilic acid	Meclofenamate	50 to 100 mg four times daily	400	\$220 for 50-mg dose
Cyclooxygenase-2 inhibitor	Celecoxib (Celebrex)	200 mg two times daily	400	\$240
Nonacidic agent	Nabumetone	1,000 to 2,000 mg one or two times daily	2,000	\$77 to \$98 (generic) and \$107 (brand) for 1,000 mg once daily

*—Estimated cost to the pharmacist based on average wholesale prices (rounded to the nearest dollar) in Red Book. Montvale, N.J.: Medical Economics Data; 2008. Cost to the patient will be higher, depending on prescription filling fee. Cost is based on standard dosage unless otherwise indicated. Some of these medications are available at considerable savings through local and national pharmacy discount programs.

useful addition to analgesic therapy.²¹ Gabapentin (Neurontin) may provide short-term relief in patients with radiculopathy.²

NONPHARMACOLOGIC TREATMENT OPTIONS

Patients commonly use nonpharmacologic treatment options, with or without consulting a physician.

Forty-five percent of patients with low back pain see a chiropractor, 24 percent use massage, 11 percent get acupuncture, and 7 percent try meditation.²²

Acupuncture provides short-term relief of chronic low back pain, improves functioning, and works as an adjunct to other therapeutic options. It has not been shown to be more effective than other treatments.^{23,24}

SORT: KEY RECOMMENDATIONS FOR PRACTICE

<i>Clinical recommendation</i>	<i>Evidence rating</i>	<i>References</i>
Acetaminophen and NSAIDs are first-line medications for treating chronic low back pain.	A	2, 14
Imaging, such as lumbar spine radiography, should be delayed at least one to two months in patients with nonspecific low back pain without red flags for serious disease.	C	6
Evaluation of psychosocial problems and “yellow flags” are useful in identifying patients with chronic low back pain who have a poor prognosis.	B	8, 9
Treatment options		
<i>Beneficial</i>		
Analgesics (acetaminophen, tramadol [Ultram])	A	2, 15-17
NSAIDs	A	2, 14, 17
Acupuncture	A	2, 22-24
Multidisciplinary rehabilitation	A	2, 27, 28
<i>Likely to be beneficial</i>		
Herbal medications (devil’s claw, white willow bark, topical cayenne)	B	18
Tricyclic antidepressants	B	2, 21
Exercise therapy	B	2, 25, 26
Behavior therapy	B	2
Massage	B	2, 29
Spinal manipulation	B	2, 30, 31
<i>Trade-off between benefit and harm</i>		
Muscle relaxants (short-term use)	B	17
Opioids	B	2, 19
<i>Insufficient or conflicting data</i>		
Antiepileptic medication (gabapentin [Neurontin]) for radicular symptoms	C	2
Viniyoga	C	2
Back school	C	35
Low-level laser therapy	C	2
Lumbar supports	C	2
Prolotherapy	C	34
Short wave diathermy	C	2
Traction	C	2, 33
Transcutaneous electrical nerve stimulation	C	2
Ultrasound	C	2
Epidural steroid injection	C	8, 36, 37

NSAIDs = nonsteroidal anti-inflammatory drugs.

A = consistent, good-quality patient-oriented evidence; B = inconsistent or limited-quality patient-oriented evidence; C = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, go to <http://www.aafp.org/afpsort.xml>.

clinically important and statistically significant differences between intervention and control groups.^{25,26}

Behavior therapy is as effective as exercise therapy for short-term relief of chronic low back pain. Consistent evidence supports cognitive behavior therapy and progressive relaxation for short-term improvement, whereas biofeedback techniques have produced mixed results. Combining behavior therapy with other modalities does not have an additive effect.²

Multidisciplinary rehabilitation programs that include a physician and at least one additional intervention (psychological, social, or vocational) alleviate subjective disability, reduce pain, return persons to work five weeks earlier, and after returning to work, reduce the amount of sick time in the first year by seven days. Benefits persist for up to five years.^{27,28}

Acupuncture massage and pressure point massage are mildly helpful for reducing chronic low back pain, and the benefits last for up to one year. Massage appears to be most effective when combined with exercise, stretching, and education.²⁹

Spinal manipulation provides modest short- and long-term relief of back pain, improves psychological well-being, and increases functioning.^{2,30} The benefits derived are not dependent on the type of training of the manipulator because osteopathic and chiropractic outcomes appear to be similar.³¹

One therapeutically directed style of yoga (Viniyoga) may provide some relief of chronic back pain. Six weeks of yoga decreased medication use and provided more pain relief than exercise and self-care. Other forms of yoga have mixed results in small studies, and at this time there is not enough evidence to recommend them.³²

Fifty-one to 64 percent of patients are willing to try acupuncture if recommended by their physician.²²

Exercise therapy, focusing on strengthening and stabilizing the core muscle groups of the abdomen and back, appears to produce small improvements in pain and functioning in patients with chronic low back pain. However, few studies (i.e., six of the 43 studies included in a Cochrane review) have been able to demonstrate

Back schools, low-level laser therapy, lumbar supports, prolotherapy, short wave diathermy, traction, transcutaneous electrical nerve stimulation, and ultrasound have negative or conflicting evidence of effectiveness.³²⁻³⁵

EPIDURAL STEROID INJECTIONS

Epidural steroid injections may help patients with radicular symptoms. Studies have found conflicting results,

but the trend is toward a small improvement for up to three months after injection.³⁶ There is no evidence to support the use of epidural steroid injections in patients without radicular symptoms,³⁷ and injections are less effective in patients with severe spinal stenosis and those with stenotic lesions encompassing more than three lumbar levels.^{37,38}

SURGERY

Most patients with back pain will not benefit from surgery. However, if anatomic abnormalities consistent with the distribution of pain are identified, surgery can be considered in persons who have experienced significant functional disabilities and in those with unremitting pain, especially pain lasting longer than 12 months despite multiple nonsurgical treatments. Good evidence supports the use of spinal fusion for treating back pain caused by fractures, infections, progressive deformity, or instability with spondylolisthesis.⁷ Spinal decompression, nerve root decompression, and spinal fusion have been extensively evaluated for the treatment of degenerative disorders of the spine, mostly with short-term outcomes, yielding conflicting results and questionable patient benefit.³⁹ Disk arthroplasty (replacing the original intervertebral disk with an artificial one) appears to be as effective as lumbar fusion for short-term relief of chronic low back pain, but there is no evidence of long-term relief, and concerns exist regarding the durability of the artificial disks. Intradiscal electrothermal therapy is a technique that applies heat to a damaged disk through a catheter, causing collagen contraction for structural support and ablating nearby pain-sensing nerves for pain reduction. It has been shown to provide modest pain relief, but little functional improvement.⁴⁰

REFERRAL

Referral to a pain management specialist is appropriate for patients who continue to experience severe functional impairment or unremitting pain, or when patients or physicians feel that progress has stopped or want a second opinion. In the absence of evidence to define the indications and timing of referral, a decision to refer should be left to the discretion of the physician and patient.²

The Authors

ALLEN R. LAST, MD, MPH, is program director at the Racine Family Medicine Residency Program at the Medical College of Wisconsin. He received his medical degree from the University of Wisconsin School of Medicine and Public Health, Madison, and completed a residency and a faculty development fellowship at the University of Pittsburgh (Pa.) Medical

Center St. Margaret Family Medicine Residency Program. Dr. Last received a master of public health degree from the University of Pittsburgh Graduate School of Public Health.

KAREN HULBERT, MD, is a predoctoral coordinator and an assistant professor at the Racine Family Medicine Residency Program at the Medical College of Wisconsin. She received her medical degree from Rosalind Franklin University of Medicine and Science, Chicago (Ill.) Medical School, and completed a family medicine residency at St. Paul (Minn.) Ramsey Medical Center.

Address correspondence to Allen R. Last, MD, MPH, Medical College of Wisconsin, 1320 Wisconsin Ave., Racine, WI 53403 (e-mail: alast@mcw.edu). Reprints are not available from the authors.

Author disclosure: Nothing to disclose.

REFERENCES

- Haldeman S, Dagenais S. A supermarket approach to the evidence-informed management of chronic low back pain. *Spine J*. 2008;8(1):1-7.
- Chou R, Qaseem A, Snow V, et al., for the Clinical Efficacy Assessment Subcommittee of the American College of Physicians, American Pain Society Low Back Pain Guidelines Panel. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society [published correction appears in *Ann Intern Med*. 2008;148(3):247-248]. *Ann Intern Med*. 2007;147(7):478-491.
- Deyo RA, Weinstein JN. Low back pain. *N Engl J Med*. 2001;344(5):363-370.
- Devillé WL, van der Windt DA, Dzaferagic A, Bezemer PD, Bouter LM. The test of Lasègue: systematic review of the accuracy in diagnosing herniated discs. *Spine*. 2000;25(9):1140-1147.
- Kinkade S. Evaluation and treatment of acute low back pain. *Am Fam Physician*. 2007;75(8):1181-1188.
- Bradley WG Jr, Seidenwurm DJ, Brunberg JA, et al. Expert Panel on Neurologic Imaging. Low back pain. American College of Radiology; 2005. http://www.acr.org/SecondaryMainMenuCategories/quality_safety/app_criteria/pdf/ExpertPanelonNeurologicImaging/LowBackPainDoc7.aspx. Accessed March 24, 2009.
- Don AS, Carragee E. A brief overview of evidence-informed management of chronic low back pain with surgery. *Spine J*. 2008;8(1):258-265.
- vanWijk RM, Geurts JW, Lousberg R, et al. Psychological predictors of substantial pain reduction after minimally invasive radiofrequency and injection treatments for chronic low back pain. *Pain Med*. 2008;9(2):212-221.
- New Zealand Guidelines Group. Acute Low Back Pain Guide. Wellington, New Zealand; October 2004. http://www.nzgg.org.nz/guidelines/0072/acc1038_col.pdf. Accessed March 24, 2009.
- Jensen S. Back pain—clinical assessment. *Aust Fam Physician*. 2004;33(6):393-401.
- Mallen CD, Peat G, Thomas E, Dunn KM, Croft PR. Prognostic factors for musculoskeletal pain in primary care: a systematic review. *Br J Gen Pract*. 2007;57(541):655-661.
- Yelland MJ, Schluter PJ. Defining worthwhile and desired responses to treatment of chronic low back pain. *Pain Med*. 2006;7(1):38-45.
- Gourlay DL, Heit HA, Almahrezi A. Universal precautions in pain medicine: a rational approach to the treatment of chronic pain. *Pain Med*. 2005;6(2):107-112.
- Roelofs PD, Deyo RA, Koes BW, Scholten RJ, van Tulder MW. Non-steroidal anti-inflammatory drugs for low back pain. *Cochrane Database Syst Rev*. 2008;(1):CD000396.
- Schnitzer TJ, Gray WL, Paster RZ, Kamin M. Efficacy of tramadol in treatment of chronic low back pain. *J Rheumatol*. 2000;27(3):772-778.

Low Back Pain

16. Ruoff GE, Rosenthal N, Jordan D, Karim R, Kamin M, for the Protocol CAPSS-112 Study Group. Tramadol/acetaminophen combination tablets for the treatment of chronic lower back pain: a multicenter, randomized, double-blind, placebo-controlled outpatient study. *Clin Ther*. 2003; 25(4):1123-1141.
17. Malanga G, Wolff E. Evidence-informed management of chronic low back pain with nonsteroidal anti-inflammatory drugs, muscle relaxants, and simple analgesics. *Spine J*. 2008;8(1):173-184.
18. Gagnier JJ, van Tulder M, Berman B, Bombardier C. Herbal medicine for low back pain. *Cochrane Database Syst Rev*. 2006;(2):CD004504.
19. Deshpande A, Furlan A, Mailis-Gagnon A, Atlas S, Turk D. Opioids for chronic low-back pain. *Cochrane Database Syst Rev*. 2007;(3):CD004959.
20. Chang G, Chen L, Mao J. Opioid tolerance and hyperalgesia. *Med Clin North Am*. 2007;91(2):199-211.
21. Staiger TO, Gaster B, Sullivan MD, Deyo RA. Systematic review of antidepressants in the treatment of chronic low back pain. *Spine*. 2003;28(22):2540-2545.
22. Sherman KJ, Cherkin DC, Connelly MT, et al. Complementary and alternative medical therapies for chronic low back pain: What treatments are patients willing to try? *BMC Complement Altern Med*. 2004;4:9.
23. Furlan AD, van Tulder MW, Cherkin DC, et al. Acupuncture and dry-needling for low back pain. *Cochrane Database Syst Rev*. 2005;(1):CD001351.
24. Witt CM, Jena S, Selim D, et al. Pragmatic randomized trial evaluating the clinical and economic effectiveness of acupuncture for chronic low back pain. *Am J Epidemiol*. 2006;164(5):487-496.
25. Hayden JA, van Tulder MW, Malmivaara A, Koes BW. Exercise therapy for treatment of non-specific low back pain. *Cochrane Database Syst Rev*. 2005;(3):CD000335.
26. van Tulder M, Malmivaara A, Hayden J, Koes B. Statistical significance versus clinical importance: trials on exercise therapy for chronic low back pain as example. *Spine*. 2007;32(16):1785-1790.
27. Karjalainen K, Malmivaara A, van Tulder M, et al. Multidisciplinary biopsychosocial rehabilitation for subacute low back pain among working age adults. *Cochrane Database Syst Rev*. 2003;(2):CD002193.
28. Vollenbroek-Hutten MM, Hermens HJ, Wever D, Gorter M, Rinkel J, Ijzerman MJ. Differences in outcome of a multidisciplinary treatment between subgroups of chronic low back pain patients defined using two multiaxial assessment instruments: the multidimensional pain inventory and lumbar dynamometry. *Clin Rehabil*. 2004;18(5):566-579.
29. Furlan AD, Imamura M, Dryden T, Irvin E. Massage for low-back pain. *Cochrane Database Syst Rev*. 2008;(4):CD001929.
30. Williams NH, Hendry M, Lewis R, Russell I, Westmoreland A, Wilkinson C. Psychological response in spinal manipulation (PRISM): a systematic review of psychological outcomes in randomised controlled trials. *Complement Ther Med*. 2007;15(4):271-283.
31. Assendelft WJ, Morton SC, Yu EI, Suttrop MJ, Shekelle PG. Spinal manipulative therapy for low back pain. *Cochrane Database Syst Rev*. 2004;(1):CD000447.
32. Chou R, Huffman LH, for the American Pain Society and the American College of Physicians. Nonpharmacologic therapies for acute and chronic low back pain: a review of the evidence for an American Pain Society/American College of Physicians clinical practice guideline [published correction appears in *Ann Intern Med*. 2008;148(3):247-248]. *Ann Intern Med*. 2007;147(7):492-504.
33. Clarke JA, van Tulder MW, Blomberg SE, et al. Traction for low-back pain with or without sciatica. *Cochrane Database Syst Rev*. 2007;(2):CD003010.
34. Dagenais S, Yelland MJ, Del Mar C, Schoene ML. Prolotherapy injections for chronic low-back pain. *Cochrane Database Syst Rev*. 2007;(2):CD004059.
35. Engers A, Jellema P, Wensing M, van der Windt DA, Grol R, van Tulder MW. Individual patient education for low back pain. *Cochrane Database Syst Rev*. 2008;(1):CD004057.
36. Armon C, Argoff CE, Samuels J, Backonja MM, for the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. Assessment: use of epidural steroid injections to treat radicular lumbosacral pain. *Neurology*. 2007;68(10):723-729.
37. DePalma MJ, Slipman CW. Evidence-informed management of chronic low back pain with epidural steroid injections. *Spine J*. 2008;8(1):45-55.
38. Kapural L, Mekhail N, Bena J, et al. Value of the magnetic resonance imaging in patients with painful lumbar spinal stenosis (LSS) undergoing lumbar epidural steroid injections. *Clin J Pain*. 2007;23(7):571-575.
39. Gibson JN, Waddell G. Surgery for degenerative lumbar spondylosis. *Cochrane Database Syst Rev*. 2005;(4):CD001352.
40. Derby R, Baker RM, Lee CH, Anderson PA. Evidence informed management of chronic low back pain with intradiscal electrothermal therapy. *Spine J*. 2008;8(1):80-95.

Figure. Summary of the American College of Physicians guideline on noninvasive treatments for acute, subacute, or chronic low back pain.



Summary of the American College of Physicians Guideline on Noninvasive Treatments for Acute, Subacute, or Chronic Low Back Pain

Disease/Condition	Low back pain
Target Audience	All clinicians
Target Patient Population	Adults with acute, subacute, or chronic low back pain
Interventions Evaluated	<p>Pharmacologic interventions: NSAIDs, nonopioid analgesics, opioid analgesics, tramadol and tapentadol, antidepressants, SMRs, benzodiazepines, corticosteroids, antiepileptic drugs</p> <p>Nonpharmacologic interventions: interdisciplinary or multicomponent rehabilitation; psychological therapies; exercise and related interventions, such as yoga or tai chi; complementary and alternative medicine therapies, including spinal manipulation, acupuncture, and massage; passive physical modalities, such as heat, cold, ultrasound, transcutaneous electrical nerve stimulation, electrical muscle stimulation, interferential therapy, short-wave diathermy, traction, LLLT, lumbar supports/braces</p>
Outcomes Evaluated	Pain, function, health-related quality of life, work disability/return to work, global improvement, number of back pain episodes or time between episodes, patient satisfaction, adverse effects
Benefits	<p>Acute low back pain</p> <p>Pharmacologic</p> <ul style="list-style-type: none"> NSAIDs: improved pain and function (small effect) SMRs: improved pain (small effect) <p>Nonpharmacologic</p> <ul style="list-style-type: none"> Heat wrap: improved pain and function (moderate effect) Massage: improved pain and function (at 1 but not 5 wk) (small to moderate effect) Acupuncture: improved pain (small effect) Spinal manipulation: improved function (small effect) <p>Chronic low back pain</p> <p>Pharmacologic</p> <ul style="list-style-type: none"> NSAIDs: improved pain (small to moderate effect) and function (no to small effect) Opioids: improved pain and function (small effect) <ul style="list-style-type: none"> Tramadol: improved pain (moderate effect) and function (small effect) Buprenorphine (patch or sublingual): improved pain (small effect) Duloxetine: improved pain and function (small effect) <p>Nonpharmacologic</p> <ul style="list-style-type: none"> Exercise: improved pain and function (small effect) Motor control exercise: improved pain (moderate effect) and function (small effect) Tai chi: improved pain (moderate effect) and function (small effect) Mindfulness-based stress reduction: improved pain and function (small effect) Yoga: improved pain and function (small to moderate effect, depending on comparator) Progressive relaxation: improved pain and function (moderate effect) Multidisciplinary rehabilitation: improved pain (moderate effect) and function (no to small effect) Acupuncture: improved pain (moderate effect) and function (no to moderate effect, depending on comparator) LLLT: improved pain and function (small effect) Electromyography biofeedback: improved pain (moderate effect) Operant therapy: improved pain (small effect) Cognitive behavioral therapy: improved pain (moderate effect) Spinal manipulation: improved pain (small effect) <p>Radicular low back pain</p> <ul style="list-style-type: none"> Exercise: improved pain or function (small effect)
Harms	<p>Generally poorly reported</p> <p>Pharmacologic</p> <ul style="list-style-type: none"> NSAIDs: increased adverse effects compared with placebo and acetaminophen (COX-2–selective NSAIDs decreased risk for adverse effects compared with traditional NSAIDs) Opioids: nausea, dizziness, constipation, vomiting, somnolence, and dry mouth SMRs: increased risk for any adverse event and central nervous system adverse events (mostly sedation) Benzodiazepines: somnolence, fatigue, lightheadedness Antidepressants: increased risk for any adverse event <p>Nonpharmacologic</p> <ul style="list-style-type: none"> Poorly reported, but no increase in serious adverse effects

Continued on following page

<p>Recommendations</p>	<p>Recommendation 1: Given that most patients with acute or subacute low back pain improve over time regardless of treatment, clinicians and patients should select nonpharmacologic treatment with superficial heat (moderate-quality evidence), massage, acupuncture, or spinal manipulation (low-quality evidence). If pharmacologic treatment is desired, clinicians and patients should select nonsteroidal anti-inflammatory drugs or skeletal muscle relaxants (moderate-quality evidence). (Grade: strong recommendation)</p> <p>Recommendation 2: For patients with chronic low back pain, clinicians and patients should initially select nonpharmacologic treatment with exercise, multidisciplinary rehabilitation, acupuncture, mindfulness-based stress reduction (moderate-quality evidence), tai chi, yoga, motor control exercise, progressive relaxation, electromyography biofeedback, low-level laser therapy, operant therapy, cognitive behavioral therapy, or spinal manipulation (low-quality evidence). (Grade: strong recommendation)</p> <p>Recommendation 3: In patients with chronic low back pain who have had an inadequate response to nonpharmacologic therapy, clinicians and patients should consider pharmacologic treatment with nonsteroidal anti-inflammatory drugs as first-line therapy, or tramadol or duloxetine as second-line therapy. Clinicians should only consider opioids as an option in patients who have failed the aforementioned treatments and only if the potential benefits outweigh the risks for individual patients and after a discussion of known risks and realistic benefits with patients. (Grade: weak recommendation, moderate-quality evidence)</p>
<p>High-Value Care</p>	<p>Clinicians should reassure patients that acute or subacute low back pain usually improves over time regardless of treatment and should avoid prescribing costly and potentially harmful treatments. Systemic steroids were not shown to provide benefit and should not be prescribed for patients with acute or subacute low back pain, even with radicular symptoms. For treatment of chronic low back pain, clinicians should select therapies that have the fewest harms and lowest costs. Clinicians should avoid prescribing costly therapies and those with substantial potential harms, such as long-term opioids, and pharmacologic therapies that were not shown to be effective, such as tricyclic antidepressants and selective serotonin reuptake inhibitors.</p>
<p>Clinical Considerations</p>	<p>Clinicians should inform patients with acute or subacute low back pain of the generally very favorable outcome. Thus, patients can avoid potentially harmful and costly tests and treatments.</p> <p>Clinicians should advise patients with acute, subacute, or chronic low back pain to remain active as tolerated.</p> <p>Improvements in pain and function due to pharmacologic and nonpharmacologic interventions were small and often showed no clear differences compared with controls.</p> <p>Few differences in recommended therapies were found when they were studied in head-to-head trials. Therefore, clinicians should base treatment recommendations on patient preferences that also minimize harms and costs.</p>

COX-2 = cyclooxygenase-2; LLLT = low-level laser therapy; NSAID = nonsteroidal anti-inflammatory drug; SMR = skeletal muscle relaxant.

Note Reprinted from “Noninvasive Treatments for Acute, Subacute and Chronic Low Back Pain: A Clinical Practice Guideline From the American College of Physicians”, by Qaseem A (February 2017) Retrieved from <http://annals.org/aim/article/2603231/noninvasive-treatments-acute-subacute-chronic-low-back-pain>

References

1. Qaseem A, Wilt TJ, McLean RM, Forcica MA, for the Clinical Guidelines Committee of the American College of Physicians. Noninvasive Treatments for Acute, Subacute, and Chronic Low Back Pain: A Clinical Practice Guideline from The American College of Physicians. *Ann Intern Med.* 2017; 166:514–530. Retrieved from <http://annals.org/aim/article/2603231/noninvasive-treatments-acute-subacute-chronic-low-back-pain>.