

OPTIMA FAMILY CARE MEDALLION 4.0

PHARMACY/MEDICAL PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: *The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; fax to 1-844-723-2094. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. Incomplete form will delay authorization process.*

Drug Requested: **Botulinum Toxin Injections®, Type A (Medical)**

Botox® (onabotulinumtoxinA) (J0585)

Xeomin® (incobotulinumtoxinA) (J0588)

DRUG INFORMATION: *Check the applicable box above and complete all information below or authorization process will be delayed.*

Drug Name/Form/Strength/Quantity: _____

Dosing Schedule: _____ **Length of Therapy:** _____

Diagnosis: _____ **ICD Code:** _____

- **Max quantity limits: 400 units in a 3-month period**
- **Cosmetic indications are excluded.**

CLINICAL CRITERIA: *Check **one** of the diagnoses below. Applicable lines **MUST** be checked to ensure the authorization process will **NOT** be delayed.*

**** Medical notes must be submitted to support each line checked on this request. ****

- | | |
|---|--|
| <ul style="list-style-type: none"> <input type="checkbox"/> Achalasia, Primary idiopathic esophageal <ul style="list-style-type: none"> <input type="checkbox"/> The patient has failed or had a clinically significant adverse reaction to conventional therapy (nitrates or calcium channel blockers) <p style="text-align: center;">OR</p> <input type="checkbox"/> The patient ineligible for surgical treatment due to advance age or multiple co-morbidities (poor surgical risk) <p style="text-align: center;">OR</p> <ul style="list-style-type: none"> <input type="checkbox"/> The patient is at high risk of complications of pneumatic dilation or surgical myotome <p style="text-align: center;">OR</p> <ul style="list-style-type: none"> <input type="checkbox"/> Failure of prior myotomy or dilation <p style="text-align: center;">OR</p> <ul style="list-style-type: none"> <input type="checkbox"/> The patient has an epiphrenic diverticulum or hiatal hernia, both of which increase the risk of dilation-induced perforation | <ul style="list-style-type: none"> <input type="checkbox"/> Cervical Dystonia (spasmodic torticollis) and Mixed Cervical Dystonia <input type="checkbox"/> Chronic Migraine Headache Prophylaxis
Patients must have met ALL the following criteria: <ul style="list-style-type: none"> <input type="checkbox"/> Headaches ≥ 15 days/month <input type="checkbox"/> Headaches last ≥ 4 hours/day <input type="checkbox"/> Current use of at least one migraine prophylaxis drug <input type="checkbox"/> Predominant rescue medication is NOT an opioid <input type="checkbox"/> CVA-related spasticity within 1 year of onset <input type="checkbox"/> Drooling in Parkinson's Disease <input type="checkbox"/> Essential hand tremor in patients who fail oral agents <input type="checkbox"/> Hand Dystonia <input type="checkbox"/> Hemifacial spasm <input type="checkbox"/> Hirschsprung's Disease <input type="checkbox"/> Laryngeal Dysphonia – Spastic <input type="checkbox"/> Laryngeal Dystonia (adductor spasmodic dysphonia) <input type="checkbox"/> Laryngeal Spasm <input type="checkbox"/> Motor tics <input type="checkbox"/> Neurogenic detrussor overactivity and/or detrussor sphincter dyssynergia <input type="checkbox"/> Orofacial Dyskinesia |
| <ul style="list-style-type: none"> <input type="checkbox"/> Achalasia, Internal anal sphincter (IAS) <ul style="list-style-type: none"> <input type="checkbox"/> Patient has not responded to treatment with laxatives <p style="text-align: center;">AND</p> <input type="checkbox"/> Patient has not responded to or is not a candidate for anal sphincter myectomy <input type="checkbox"/> Anal Fissure – Chronic <ul style="list-style-type: none"> <input type="checkbox"/> The patient has failed (at least 60 days) topical nitroglycerin or topical calcium channel blocker <input type="checkbox"/> Blepharospasm <input type="checkbox"/> Cerebral Palsy – Dynamic Contracture <input type="checkbox"/> Cerebral Palsy – Spasticity (including diplegia, hemiplegia, paraplegia, or quadriplegia) | |

(continued on next page)

Overactive Bladder

Patients must have met ALL the following criteria:

- A diagnosis of incontinence
- Symptoms of urge incontinence, urgency, and frequency (experienced at least 3 urinary incontinence episodes and at least 24 micturitions in 3 days)
- 8-12 week trial and failure of behavioral therapy (e.g. bladder training, control strategies, pelvic floor muscle training, fluid management)
- Failed or inadequate response to anticholinergic therapy within the last 9 months (4-8 week trial per agent)

- 2 anticholinergic agents and 1 β -3 adenosine receptor agonist (*will require PA*); or

- 1 anticholinergic agent and 1 alpha blocker and 1 β -3 adenosine receptor agonist (*will require PA*)

Please indicate drugs used: _____

- Strabismus** (injections done in lieu of coverage for surgery)
- Synkinetic Eyelid Closure – VII Cranial Nerve**
- Torticollis**

Medication being provided by (check applicable box below):

- Physician's office** **OR** **Specialty Pharmacy - PropriumRx**

****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.****

Previous therapies will be verified through pharmacy paid claims or submitted chart notes.

Patient Name: _____

Member Optima #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

REVISED/UPDATED: 7/1/2018