

OPTIMA HEALTH COMMUNITY CARE

PHARMACY PRIOR AUTHORIZATION/STEP-EDIT REQUEST*

Directions: The prescribing physician must sign and clearly print name (preprinted stamps not valid) on this request. All other information may be filled in by office staff; fax to 1-800-319-5003. No additional phone calls will be necessary if all information (including phone and fax #s) on this form is correct. Incomplete form will delay authorization process.

Drug Requested:

Repatha™ (evolocumab) (Non-Preferred)

Medicaid

DRUG INFORMATION: Complete information below. If incomplete, authorization process will be delayed.

Drug Form/Strength: _____

Dosage Schedule: _____ **Length of Therapy:** _____

Length of Authorizations: Initial Approval – three (3) months; Renewal Approval – six (6) months)

Quantity Limit:

ASCVD or HeFH: 2 pens or syringes per month

HoFH: 3 pens or syringes per month

CLINICAL CRITERIA: The following criteria must be met to qualify or authorization process will be delayed.

Initial Criteria Approval – Three (3) months

- Age \geq 18 years if diagnosis is:
 - atherosclerotic cardiovascular disease (ASCVD); **OR**
 - heterozygous familial hypercholesterolemia (HeFH); **OR**
- Age \geq 13 years if diagnosed with homozygous familial hypercholesterolemia (HoFH); **AND**
- Prescribed by or in consultation with a specialist (including cardiologists, lipidologists, or endocrinologists); **AND**
- Diagnosis of ASCVD, HeFH as confirmed by genotyping or by clinical criteria (“definite FH” using either the Simon Broome or WHO/Dutch Lipid Network criteria), or HoFH as confirmed by either:
 - Documented DNA test for functional mutation(s) in both LDL receptor alleles or alleles known to affect LDL receptor functionality **OR**
 - A history of an untreated LDL-C concentration >500 mg/dL and triglycerides <300 mg/dL and both parents with documented untreated TC >250 mg/dL; **AND**
- Prior treatment history with highest available dose or maximally-tolerated dose of high intensity statin (atorvastatin or rosuvastatin) **AND** ezetimibe for at least three continuous months with failure to reach target LDL-C (70 mg/dL for patients with clinical ASCVD and 100 mg/dL for patients with HeFH or HoFH and no history of clinical ASCVD)
- If the patient is not able to use a maximum dose of atorvastatin or rosuvastatin due to muscle symptoms, documentation of a causal relationship must be established between statin use and muscle symptoms. Documentation must demonstrate that the patient experienced pain, tenderness, stiffness, cramping, weakness, and/or fatigue and all of the following:
 - Muscle symptoms resolve after discontinuation of statin; **AND**
 - Muscle symptoms occurred when re-challenged at a lower dose of the same statin; **AND**
 - Muscle symptoms occurred after switching to an alternative statin; **AND**
 - Documentation ruling out non-statin causes of muscle symptoms (e.g., hypothyroidism, reduced renal function, reduced hepatic function, rheumatologic disorders, such as polymyalgia rheumatica, steroid myopathy, vitamin D deficiency, or primary muscle disease); **OR**
- The patient has been diagnosed with statin-induced rhabdomyolysis
 - The diagnosis should be supported by acute neuromuscular illness or dark urine; **AND**
 - an acute elevation in creatine kinase (usually $>5,000$ IU/L or five times the upper limit or normal)
- If the patient failed to reach target LDL-C (<70 mg/dL for patients with clinical ASCVD and <100 mg/dL for patients with HeFH or HoFH and no history of clinical ASCVD), adherence to maximally tolerated statin and ezetimibe has been verified using pharmacy claims data and the patient is determined to be compliant for at least three consecutive months prior to the lipid panel demonstrating suboptimal reduction.
- Maximally-tolerated statin will continue to be used in conjunction with evolocumab; **AND**
- Patient has not had a prior trial and failure of an alternative PCSK9 inhibitor.

(continued on next page)

Renewal Criteria – Approval for 6 months (May be requested by PCP):

- Lipid panel showing a further reduction in LDL-C compared to the labs prior to initiating evolocumab; **AND**
- Continued adherence to maximally-tolerated statin dose established prior to the original evolocumab approval

Use of samples to initiate therapy **does not meet step-edit/preauthorization criteria.**

****Previous therapies will be verified through pharmacy paid claims or submitted chart notes.****

Patient Name: _____

Member Optima #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

REVISED/UPDATED: ~~6/29/2017~~ 8/31/2017.