

Cirrhosis requires 2 liver assessments with Lab values & symptoms correlating with Cirrhosis.

Submit a Liver assessment documenting stage 3 or stage 4 hepatic fibrosis including one (1) of the following:

Please Note: Contra-Indication to a liver assessment would lead to a denial.

- Liver biopsy confirming:
 - METAVIR score of F3 - F4 Knodell fibrosis score (last #-reported separately) F 3-4
 - Ishak stage score of F4 –F6 Batts-Ludwig stage 3 -4
- Transient elastography (FibroScan) score greater than or equal to 9.5 kPa
- FibroTest (FibroSure) score of greater ≥ 0.59 (F3) or ≥ 0.75 (F4)
- Shear wave elastography (ElastPQ) score of 12.0-21.0+kPa or 2.00-2.64+m/s
- Shear wave (SWE supersonic tech) score of greater ≥ 8.7 kPa (1.70 m/s) (F3) OR 10.4kPa (1.86m/s) (F4)
- Shear wave (VTTQ) Siemens score of greater ≥ 1.55 m/s (F3) OR 1.80m/s(F4)

Genotype 1, 2, 3, 4, 5, or 6 HCV	
Patient Population	Recommended Treatment Regimen
Patients without cirrhosis and patients with compensated cirrhosis (Child Pugh A)	EPCLUSA for 12 weeks
Patients with decompensated cirrhosis (Child Pugh B and C)	EPCLUSA + ribavirin for 12 weeks

****Use of samples to initiate therapy does not meet step edit/ preauthorization criteria.****

Previous therapies will be verified through pharmacy paid claims or submitted chart notes.

Patient Name: _____

Member Optima #: _____ Date of Birth: _____

Prescriber Name: _____

Prescriber Signature: _____ Date: _____

Office Contact Name: _____

Phone Number: _____ Fax Number: _____

DEA OR NPI #: _____

*Approved by Pharmacy and Therapeutics Committee: 7/21/2016

REVISED/UPDATED: 8/3/2016; 9/20/2016; 12/16/2016 8/12/2017;